

Table 1. Comparison of unprocessed vs. chemically processed crawfish waste meal.

Nutrient	*Control	*Treatment #1	*Treatment #2	*Treatment #3
Crude Protein (%)	^b 31.40	^a 61.10	^a 57.90	^a 61.60
Crude Fiber (%)	^b 13.00	^a 33.00	^a 31.50	^a 29.60
Calcium (%)	^a 15.40	^b .80	^c .38	^d .03
Phosphorous (%)	^a 1.36	^b .36	^c .25	^d .07
Ca:P Ratio	^a 11.30	^b 2.20	^c 1.57	^d .43

a,b,c,d

Superscripts denote statistical significant differences at ($P < .01$).

*Control denotes no chemical treatment of the crawfish waste meal. Treatment #1 denotes the use of a strong acid (three molar concentration and two rinses of the product). Treatment #2 denotes the use of a strong acid (three molar concentration and three rinses of the product). Treatment #3 denotes the use of a strong acid (four molar concentration and two rinses of the product).

0992796622660

Table 2. Comparison of lysine and methionine profiles of unprocessed crawfish waste meal, soybean meal and chemically processed crawfish waste meal.

Nutrient	*Control	*Soybean Meal	*Processed Crawfish Meal
Crude Protein (%)	^a 34.60%	^b 46.80%	^c 61.30%
Lysine (gm/100 gm)	^a 1.34%	^b 2.90	^b 2.30
Methionine (gm/100 gm)	^a .53	^a .57	^b .94
Calcium (%)	^a 14.90%	^b .38	^c .49
Phosphorous (%)	^a 1.50	^b .67	^c .12


a,b,c

Superscripts denote statistical significant differences at ($P < .01$).

*Control (crawfish meal) and soybean meal samples were chemically treated. The processed crawfish waste meal was treated with a strong acid (three molar concentration and two rinses of the product).

09927996-081001

United States Patent & Trademark Office
Office of Initial Patent Examination -- Scanning Division



Application deficiencies found during scanning:

☐ Page(s) _____ of only 5 drawings were ~~not~~ present
for scanning. (Document title)

☐ Page(s) _____ of _____ were not present
for scanning. (Document title)

☐ *Scanned copy is best available.*

FOOTED 966/2660